X-ray Study of Dynamical Evolution in Clusters of Galaxies

Tae Furusho

Department of Physics, Tokyo Metropolitan University
1-1 Minami-Ohsawa, Hachioji, Tokyo 192-0397, Japan

Submitted to Tokyo Metropolitan University
for the Degree of Doctor of Philosophy (Physics)
January 2001

Thie wortk is based on analysis of data from the ISAS
satellite ASCA in collaboration with ASCA team.

Present address:
Code 662, NASA/GSFC, Greenbelt, MD 20771, USA
E-mail: furusho@olegacy.gsfc.nasa.gov

Abstract

We present two-dimensional temperature maps of four nearby bright clusters of galaxies A1060, AWM7, the Centaurus cluster, and the Perseus cluster, based on multi-pointing observations with the *ASCA* GIS. The pointing fields cover the whole cluster regions of $1 - 1.8^\circ$ corresponding to about 2–3 h_{50}^{-1} Mpc. The temperatures were derived from hardness ratios, including corrections for the effects of the XRT response. The central cool components were subtracted for the Centaurus and the Perseus clusters, and metallicity gradients in AWM7, the Centaurus and the Perseus clusters were also corrected.

The intracluster medium in A1060 and AWM7 is almost isothermal from the center to the outer regions with temperatures of 3.3 and 3.9 keV, respectively. The Centaurus cluster exhibits remarkable hot region within about $30'$ south-east from the cluster center, showing a temperature increase of +0.8 keV from the surrounding level of 3.5 keV. There are also some cool regions with lower temperatures by −1.3 keV toward the cluster edge. The temperature map of the Perseus shows more complex structure in the whole cluster region. There is an extended cool region with a diameter of $\sim 20'$ and $kT \sim 5$ keV at about $20'$ east from the cluster center. This region also shows higher surface brightness, and is surrounded by a large ring-like hot region with $kT \gtrsim 7$ keV. Another extended cool region is extending outward from the IC 310 subcluster.

The observed results indicate that A1060 and AWM7 have not experienced merging processes for recent a few Gyr. The southeast hot region near the cluster center of the Centaurus corresponds to a galaxy subcluster region from optical observations. These features suggest that a strong merger has occurred in the recent ~ 1 Gyr, and that the central cool component has survived it. For the Perseus cluster, the presence of several other hot and cool blobs suggests that this rich cluster has been formed as a result of a repetition of many subcluster mergers.

Main results of this thesis have been published as 2 papers: Furusho et al. 2001, PASJ, 53, 421 (for A1060, AWM7, and Centaurus), and Furusho et al. 2001, ApJ, 561, L165 (for Perseus)

Contents

List of Figures

List of Tables

Chapter 1

Introduction

Clusters of galaxies have been regarded as the largest virialized systems in the universe with dimensions of 1–5 Mpc across. Since the time-scale for the cluster evolution is comparable to the age of the universe, ~ 10 Gyr, their properties should reflect the initial condition and the evolutional history over the cosmological time scale.

The constituents of clusters are divided into three components: galaxies , intracluster medium (ICM), and dark matter. The galaxies are mainly observed in the optical band with the number of about a few hundreds a cluster. The ICM is a diffuse, X-ray emitting hot plasma with temperatures of typically $\times 10^{7-8}$ K. Their masses are heavier than the stellar mass by a factor of 3–5, and the distributions generally trace the gravitational potential. The dark matter occupies around 80% of the total cluster mass of $10^{14-15} M_{\odot}$. The primary energy source of the ICM is considered to be the kinetic energy released through the gas infall into the gravitational potential of the dark matter. The heating mechanism is, however, not yet fully understood. Numerical simulations based on a bottom-up formation scenario predict that shock heatings caused by mergers of smaller groups and subclusters are the important process. If mergers are playing a major role in the cluster evolution, their footprints on the ICM, in the form of morphology, temperature and metallicity, can be studied with X-ray observations.

Recent X-ray observations have shown various evidences indicating that the clusters are still evolving. Einstein and ROSAT have shown that some clusters suggest external disturbances in their irregular morphologies, and ASCA has shown two-dimensional temperature maps of nearby cluster with remarkable structures. The temperature structure persist over several Gyr even after the surface brightness distribution has been almost smoothed out with a time-scale of $\lesssim$ several hundreds Myr. Also, ASCA shows that radial profiles of metallicity in the ICM can be classified into several types that depend on such as the age, gas mixing, and cluster size. The metals are considered to be ejected from the galaxies into the ICM via galactic winds, or ram pressure stripping mainly during the starburst phase of more than several Gyr ago. These characters enable us to perform, to some extent, a time sliced study of the cluster evolution.

To reach consistent understanding of the evolutionary process of clusters, we need to look into the morphology, temperature, and metallicity structures for different richness systems in a systematic manner with enough spatial resolution. In particular, constraining the temperature structures in apparently relaxed systems that have smooth surface brightnesses is very important to know the past history in the evolution of average clusters. Since nearby clusters with $z \lesssim 0.02$ cannot be covered in the field of view of the GIS instrument on ASCA, we have to devise an analysis method to correctly handle multi-pointing data. Little effort has so far been made for such a study, mainly because of the technical difficulty. We carry out here the first attempt on the sensitive studies of the temperature and metallicity distributions for a sample of nearby clusters.

ASCA provides superior energy resolution and high sensitivity over a wide energy band 0.5–10.0 keV. These enable us to map out the temperature structure in nearby clusters with a spatial resolution of less than 100 kpc. This thesis presents temperature maps of Abell 1060, AWM7, the Centaurus cluster, and the Perseus cluster, based on the multi-pointing observations from ASCA. These clusters are all located closely to us ($z = 0.0104 - 0.0183$), emitting intense X-ray fluxes with the ICM temperatures $kT = 3 - 6$ keV. The clusters are markedly different in the strength of the central cool component, and in the shape of the metallicity profile, which would reflect different past histories of the cluster evolution. Comparison of temperature maps and metallicity distributions among the four clusters should, therefore, bring us a new important knowledge about the history of dynamical evolution as well as its connection with the central structure.

Throughout this thesis, we assume $H_0 = 50$ km s^{-1} Mpc^{-1} and $q_0 = 0.5$. The solar number abundance of Fe relative to H is taken as 4.68×10^{-5} (Anders and Grevesse 1989).

Chapter 2

The Structure and Evolution of Clusters

The hierarchical structure formation in the universe forms the basis of our study. In this chapter, we briefly review the current scenario of the structure formation, its effects on the cluster evolution, and the theoretical background. We also summarize relevant past results observed in X-rays, and previous observed properties of the hot gas and member galaxies on the four clusters studied in this thesis.

2.1 Scenario of the evolution in the universe

The evolution of the universe is divided broadly into the structure formation and the chemical evolution. The former is a dynamical process of celestial objects, and the latter represents the production of heavy elements in galaxies. Clusters of galaxies are useful probes to study both evolutions by mainly three reasons: 1) Since the dynamical time-scale of clusters is close to Hubble time, clusters retain the cosmological initial conditions well, 2) Clusters can be approximated that they consist of galaxies, hot gas, and dark matter in a spherical shape, thus they are easy to deal with, 3) Although a part of metals produced in galaxies run away into intragalactic medium, they do not escape from intracluster medium (ICM), hence we can obtain the total metal abundance without omission. In order to look into the properties of clusters, X-ray observations are particularly suited, because the X-ray emitting gas gives us the physical parameters such as temperature, mass, and metal abundance.

2.1.1 Dynamical evolution

At present, the hierarchical clustering scenario is widely supported, because it is naturally expected from the cold dark matter model. In fact, the galaxies at redshifts of ~ 5 have been observed, whereas the most distant observed clusters is at $z \sim 1$. That indicates the bottom-up scenario where small systems have been formed first. According to the scenario, large-scale structures of the universe have been formed from infinitesimally small density perturbations at the early universe through the gravitational interaction. A number of numerical simulations for the structure formation have shown producing the large-scale structures and clusters. Following an early work by White's 700-body simulations (1976), recent calculations (e.g. Eke et al. 1998) include N-body/gasdynamical simulations as shown in figure 2.1. Their results show hierarchical clustering for the hot gas and dark matter, and indicate that the clusters are formed through subcluster mergers and/or absorption of groups of galaxies.



Figure 2.1: Dot plots showing the projected particle positions in cubes of physical size 10 h^{-1} Mpc centered on a cluster at different times (Eke et al. 1998).

Figure 2.2 shows how the radii of overdense spheres behave during the expansion of the universe. For a large initial overdensity, the expansion is stopped at an early stage and the system collapses into a virialized system. A system which is now virialized must have an overdensity $\rho/\rho_{\rm crit} > 200$, and clusters of galaxies just satisfies this condition. Those which are now infalling must have $\rho/\rho_{\rm crit} > 5$ that corresponds to dense parts of superclusters of galaxies. This gives clusters of galaxies a very special place in the hierarchy of structures in the universe.



Figure 2.2: The dynamics of overdense spheres in the expanding universe. The larger initial overdensity, the earlier the sphere's expansion halts (Rees 1992).

2.1.2 Chemical evolution

The existence and distribution of the chemical elements and their isotopes are a consequence of nuclear processes that have taken place in the Big Bang, and subsequently in stars and interstellar medium. First, the gas collapses into a star by the gravitational potential of dark matter, and heavy elements produced in stars. Secondly, they are scattered into the surrounding gas through supernovae (SN). Thirdly, the gas including the heavy elements collapses into a star. The chemical evolution means a change of the total metal abundance in a galaxy through repeat of the above process.

Based on the optical observations, the element and star formation history of the universe are derived as shown in figure 2.3 (Madau 1999). The results appear an existence of a decline in the universal metal production rate at the redshift of $z < 1$ whether uncorrected or corrected for dust extinction. On the other hand, X-ray observations give the metal abundances for the elements in the ICM. Since part of metals are transfered into the ICM by galactic wind or ram pressure stripping, the total metal abundances are obtained from X-ray spectroscopy.

2.2 Observational studies of cluster evolution

Main subjects of clusters obtained from X-ray observations are morphology, temperature structure, central cool component, and metal distribution. Here we briefly summarize relevant past results from X-ray observations.



Figure 2.3: *left:* Mean comoving density of star formation as a function of cosmic time. The data points with error bars have been inferred from the UV-continuum luminosity densities. The dotted line shows the fiducial rate, $<\dot{\rho}_*>= 0.054 M_\odot$ yr^{-1} Mpc^{-3}, required to generate the observed extragalactic background light. *right:* dust corrected values ($A_{1500} = 1.2$ mag). The Hα determinations (filled triangles) together with the SCUBA lower limit (empty pentagon) have been added for comparison (Madau 1999).

2.2.1 Morphology

Figure 2.4 shows the X-ray images of A1367 and A754 in the 0.5–2.0 keV band with ROSAT PSPC. A1367 is a typical irregular cluster without a central concentration, it suggests to be a young cluster ongoing subcluster mergers. The X-ray images of A754 obviously indicates that these clusters are under a major merger. A754 also shows significant temperature structure regarding the morphology (Markevitch et al. 2000). We can expect a merging cluster based on substructures in X-ray surface brightness. However, the time-scale of apparently irregular morphology is shorter than that of temperature structure, and most of clusters show circularly symmetric surface brightness.

2.2.2 Temperature structure

Recent experiments made possible to obtain spatially resolved spectra of clusters, and are revealing the temperature structure. Markevitch et al. (1998) analyzed spatially resolved X-ray spectra of 30 nearby clusters with ASCA and ROSAT. They reported that most of them show a similar temperature decline at large radii, and about a half of the sample shows signs of merging. Determination of the dynamical structure of the clusters can constrain cosmological models because clusters should be dynamically more relaxed in a



A1367 A754

Figure 2.4: The X-ray morphology of A1367 (left) and A754 (right) several clusters of galaxies with ROSAT PSPC.

low-density universe than in a high-density one (e.g. Richstone et al. 1992). Comparing the temperature profiles with the results of cluster hydrodynamic simulations, Markevitch et al. (1998) found the low Ω ($\lesssim 0.3$) cosmological models are suitable to explain the temperature profiles.

For example, we cite two instances of 2-dimensional temperature structures of the Coma cluster. Figure 2.5 shows color-coded temperature map of the Coma cluster obtained from ASCA (Watanabe et al. 1999) and XMM-Newton (Arnaud et al. 2000). The extended north-west region is clearly hotter than the surrounding regions. The result indicates evidence of a recent merger in the Coma cluster. It is interesting to see no substructure in the X-ray image around the hot region. Hence, the temperature structure gives us much information of mergings, which is not given by the morphology only. Note that the Coma cluster also shows that the abundance distribution is almost uniform. This probably mean that the metals in the Coma cluster are mixed by mergers.

2.2.3 Central cool component

Another important feature in clusters is the existence of a cool gas at the cluster center. The presence and intensity are recognized either from central narrow peaks in the surface brightness profiles or from X-ray spectra. Their origin has often been interpreted in terms of cooling flow models that requires a multi-phase temperature structure (e.g. Fabian 1994a), whereas extensive cluster studies from ASCA spectroscopy have shown the co-existence of hot ($kT \gtrsim 3$ keV) and cool ($kT \sim 1$ keV) components in several systems (Ikebe et al. 1999). It is important to know when and how these cool components have been created in the course of cluster evolution.

If there is no strong concentration in the central region, that indicates the cluster is too young or poor to reach enough high density to start radiative cooling, or the cool com-



Figure 2.5: Color coded temperature map of the Coma cluster with ASCA (Watanabe et al. 1999) and XMM-Newton (Arnaud et al. 2000).

ponent might be disrupted by mergers. The presence of strong cool component suggests that there is no merger recently. However, when the central excess is enough strong, and subcluster is very small, it is possible to survive under a merger. It is important to know when and how these cool components have been created in the course of cluster evolution, though the expectation is very difficult.

Note that XMM Reflection Grating Spectrometers observation recently discoveries that the volume emission measure of cool component for A1795 ($z = 0.0631$), which is one of the strong cooling flow clusters, is less than a few % of the hot component at the center (Tamura et al. 2001). Also, A1835 ($z = 0.2523$) which is a strong cooling flow cluster with a $kT \sim 8$ keV shows that the emission measure of the cool gas below $kT = 2.7$ keV is much lower than expected from standard cooling flow models (Peterson et al. 2001). These results suggest that the explanation of central cool component by cooling flow model is very difficult. The detailed feature mechanism of cool component is still mysterious.

2.2.4 Metal distribution

Since heavy elements can be produced only by thermonuclear reactions in stars and supernovae, the presence of the emission lines found in the spectra from the ICM implies that heavy elements processed in galaxies largely contaminate the ICM.

Koyama et al. (1991) discovered Fe abundance increases at the center of the Virgo cluster with Ginga. From the ASCA observation, similar central abundance increases were found in several clusters (Fukazawa et al. 1994; David et al. 1996; Xu et al. 1997; Kikuchi et al. 1999). Moreover, the metal profiles are roughly divided into three types: 1) central concentration, 2) large-scale gradient, and 3) constant. For example, the metal profiles of Virgo and Centaurus cluster (Figure 2.16, Ikebe et al. 1999) are represented by the first type. They show extremely concentrated cool component associated with the cD galaxy at the cluster center. The central region is considered to survive mergers sometimes which mixed metals in the outer region. Next, a large scale gradient in the metal abundance (over 500 kpc) is detected in AWM7 (Figure 2.13, Ezawa et al. 1997) and Perseus cluster (Figure 2.4.5, Ezawa et al. 2001). These features suggest that the gas injected from individual galaxies is not efficiently mixed in the cluster space, and traces the distribution of galaxies, so that implies that the merging processes occured before starburst phase. Finally, Abell 1060 (Figure 2.10, Tamura et al. 1996) and Coma cluster show very flat metal abundance distributions. This suggests that the metals are well mixed by mergers after starburst phase. Hence, the observed metal abundance distribution enables us to study the chemical evolution related with merging history.

2.3 Theoretical background

We briefly describe here the X-ray emission mechanism, shock heating due to a merger, and characteristic time-scales as the basic theoretical background to study the cluster evolution.

2.3.1 X-ray emission from hot plasma

The ICM, having low gas density ($\sim 10^{-3}$ cm^{-3}) and a high temperature ($\sim 10^{8}$ K) radiates predominantly in the X-ray band. The emitted X-rays consist of thermal Bremsstrahlung continuum and various atomic emission lines.

The plasma volume emissivity, i.e. X-ray radiation energy emitted from a unit volume in a unit time, is given as

$$\epsilon = n_{\rm g}^2 \Lambda(T, Z). \tag{2.1}$$

The quantity $n_{\rm g}$ is the number density of the gas, and $\Lambda(T, Z)$ is the cooling function, with T and Z representing the plasma temperature and the heavy element abundance, respectively. Figure 2.6 shows the cooling function as a function of the plasma temperature assuming the cosmic abundances. The contribution of the bremsstrahlung continuum to Λ increases as $\propto T^{1/2}$. In contrast, the line contribution to Λ monotonously decreases with temperature above $T \sim 10^7$ K.

We can obtain the total X-ray luminosity by integrating equation 2.1. It is useful to define the emission integral as

$$EI = \int n_{\rm g}^2 \, dV, \tag{2.2}$$

where V is the volume of the cluster. If we assume that the ICM has a spatially-uniform temperature and abundance in the volume V, and that the ICM density is constant over the projected sky area S, then the luminosity is given as

$$L_X = \int \epsilon \, dV = EM \times S \times \Lambda(T, Z). \tag{2.3}$$

The EM is the emission measure defined as

$$EM = \int n_g^2 \, dl, \tag{2.4}$$

where l is the depth of the plasma along the line of sight. The emission integral determines the normalization of the spectrum (the overall intensity or luminosity), and the shape of the spectrum depends only on the temperature T and the heavy element abundance Z. Thus, by spectral fittings performed in the following sections, we can obtain T, Z, and EI (or EM if S is known) from the observed X-ray spectra.

2.3.2 Shock heating

Recent observations revealed statistically significant substructures in the X-ray surface brightness in many clusters (e.g. Jones & Forman 1984; Burns et al. 1994). The observed substructures in the ICM argue for remnants or ongoing merger events. The hydro/N-body simulations show that the shocks play important roles to heat up the ICM, if the merging occurs supersonically (see e.g. Ishizaka & Mineshige 1996). We briefly discuss the transition of physical conditions in the upstream to the downstream of the shock.

The mean free path of electrons (λ_e) and ions (λ_i) in a plasma are expressed as

$$\lambda_e = \lambda_i \approx 23 \left(\frac{n_e}{10^{-3}\ \mathrm{cm}^{-3}}\right)^{-1} \left(\frac{T}{10^8\ \mathrm{K}}\right)^2 \ \mathrm{kpc} \tag{2.5}$$

(Sarazin 1988, p. 156), where n_e is the number density of electrons. If the region of interest has a much larger scale than the mean free path, we may consider that the physical

Figure 2.6: Temperature dependence of the cooling function with its components for optically thin plasma containing cosmic abundances of elements (Gehrels & Williams 1993).



FIG. 1.—Temperature dependence of the cooling coefficient and its components for an optically thin plasma of cosmic abundances (Allen 1973). Line emission and thermal bremsstrahlung dominate with minor contributions from radiative recombination and two-photon continuum emission (Raymond 1988). The cooling coefficient is the volume emissivity of the gas divided by electron density and proton density. Regions of positive slope at $\lesssim 2 \times 10^5$, 10^6, 6×10^6, and $> 2.4 \times 10^7$ K (see Table 1 for exact ranges) give rise to increased thermal stability. The curve was calculated with a recently updated code from Raymond & Smith (1977; 1992, private communication), modified by us to include relativistic corrections applied to the bremsstrahlung curve. The local maximum at 8×10^6 K is known as the L-shell emission effect in laboratory plasmas with iron impurities (e.g., Griem 1983). Below $\sim 10^5$ K, the steady state equilibrium assumed in calculating this curve may not be accurate due to perturbations in the ionization balance by the heating agent.

For a system at constant pressure, the stability criterion becomes

$$\left(\frac{\partial L}{\partial T}\right)_P = \left(\frac{\partial L}{\partial T}\right)_n - \frac{n_0}{T_0}\left(\frac{\partial L}{\partial n}\right)_T < 0, \tag{3}$$

where P is pressure.

Since the radiative volume emissivity for thin plasmas varies as $T^{\alpha} n^2$ where α is roughly constant over restricted temperature ranges, the conditions for instability (assuming the heating function is temperature independent and the gas is thin to its own radiation) are $\alpha < 0$ for perturbations at constant density and $\alpha < 2$ for perturbations at constant pressure. Field (1965) argued that the constant pressure criterion is more applicable for an ideal system, but that effects such as conduction and magnetic confinement can move real plasmas toward the constant density case. It is important to note that, for either case, stability is enhanced for larger α, that is, for more positive slope in the cooling curve.

3. COOLING CURVE

The Parker-Field instability is particularly important for optically thin cosmic gases at 10^5–10^7 K, where the dominant radiation is from atomic lines. Many calculations have been performed of the radiation expected from such plasmas (Tucker & Koren 1971; Raymond, Cox, & Smith 1976; Gaetz & Salpeter 1983; Summers & McWhirter 1979; Raymond & Smith 1977; Cox & Tucker 1969; Mewe, Gronenschild, & van den Oord 1985). We have calculated, as shown in Figure 1, the total radiation output from a thin plasma using the recently updated code of Raymond & Smith (1977; 1992, private communication). The plasma is assumed to be in steady state collisional ionization equilibrium with radiative cooling balancing a temperature-independent heating source. It is assumed to have cosmic elemental abundances (Allen 1973) and high fractional hydrogen ionization. The quantity plotted is the cooling coefficient, $\Lambda(T)$, which is the volume emissivity divided by the electron density times the proton density [$\Lambda(T) \propto T^{\alpha}$ above]. Below 2×10^5 K, the dominant cooling is line emission which peaks near 10^5 K due to efficient cooling from carbon and oxygen, and then decreases toward higher temperatures as the atomic levels for the abundant low-Z elements become vacant. Above 2×10^7 K, the cooling curve increases due to thermal bremsstrahlung which we calculated including relativistic corrections.

From 2×10^5 to 2×10^7 K, the negative slope of the cooling curve ($\alpha \approx -1$) makes the gas thermally unstable. There are, however, two narrow temperature ranges of positive slope at $\sim 10^6$ and $\sim 6 \times 10^6$ K (see Fig. 1). The local maxima are caused by combinations of particular lines of Si, S, and Fe that have high emissivities over narrow temperature bands. We propose that these ranges of positive slope give rise to enhanced thermal stability. The exact ranges are given in Table 1 in addition to the well known ranges less than $\sim 2 \times 10^5$ K and greater than 2×10^7 K.

The ranges of positive slope in the cooling curve have been previously noted by Gaetz, Edgar, & Chevalier (1988) who suggested that they reduce instability in radiative shocks. Generally, the importance of the shape of the cooling curve on thermal stability has long been recognized, particularly for stellar coronae (Rosner, Tucker, & Vaiana 1978; Démoulin et al. 1987).

TABLE 1

TEMPERATURE RANGES OF POSITIVE SLOPE IN COOLING CURVE[a]

Temperature Range (K)	Energy Range (keV)	Average Temperature (K)	Dominant Emitters	Approximate α[b]
$\lesssim 2 \times 10^5$ [c]	$\lesssim 0.02$	…	C, O	3.0
$(0.79–1.4) \times 10^6$	0.068–0.12	1.1×10^6	Si, S, Fe	0.2
$(4.3–7.5) \times 10^6$	0.38–0.65	6.0×10^6	Fe	0.3
$> 2.4 \times 10^7$	> 2.1	…	Bremsstrahlung	0.4

[a] See Fig. 1 for plot of cooling curve.
[b] Cooling coefficient $\Lambda(T) \propto T^{\alpha}$. Between these temperature ranges $\alpha \approx -1$.
[c] Includes a small range from 1.0×10^5 to 1.4×10^5 K where the slope is negative.

parameters shown in figure 2.7 satisfy the 'jump' condition. It is called as Rankine-Hugoniot jump condition (e.g. Shu 1992). The mass, momentum, and energy conservation equations are expressed as

$$\rho_1 v_1 = \rho_2 v_2, \tag{2.6}$$

$$\rho_1 v_1^2 + p_1 = \rho_2 v_2^2 + p_2, \tag{2.7}$$

$$\frac{1}{2}v_1^2 + \frac{\gamma}{\gamma-1}\frac{p_1}{\rho_1} = \frac{1}{2}v_2^2 + \frac{\gamma}{\gamma-1}\frac{p_2}{\rho_2}, \tag{2.8}$$

where ρ, v, and p indicate the density, velocity, and pressure, respectively, measured from the rest frame of the shock. The suffixes of 1 and 2 represent the upstream and downstream of the shock, respectively, and γ is the ratio of specific heats. From the above equations, we find



Figure 2.7: Schematic representation of the transition region for a shock wave measured in the rest frame of the shock. Upstream gas properties are transformed to downstream ones.

$$\frac{\rho_2}{\rho_1} = \frac{(\gamma+1)\mathcal{M}_1^2}{(\gamma-1)\mathcal{M}_1^2+2}, \tag{2.9}$$

$$\frac{p_2}{p_1} = \frac{2\gamma\mathcal{M}_1^2-(\gamma-1)}{\gamma+1}, \tag{2.10}$$

$$\frac{T_2}{T_1} = \frac{p_2\cdot\rho_1}{p_1\cdot\rho_2} = \frac{[2\gamma\mathcal{M}_1^2-(\gamma-1)][(\gamma-1)\mathcal{M}_1^2+2]}{(\gamma+1)^2\mathcal{M}_1^2}, \tag{2.11}$$

where $\mathcal{M}$ indicates the Mach number.

In the limit of a weak shock, $p_1 \approx p_2$, the shock wave travels with a velocity which is close to the speed of sound, and is practically the same as an acoustic compression wave (Zel'dovich & Raizer 1966). On the other hand, in the limit of a strong shock, $p_2 \gg p_1$, $\mathcal{M}_2$ approaches a constant value $\mathcal{M}_2 \to [(\gamma-1)/2\gamma]^{1/2}$, and p_2 is expressed as

$$p_2 \approx \frac{2(\gamma-1)}{(\gamma+1)^2}\rho_2 v_1^2. \tag{2.12}$$

Thus, for the strong shock, the gas can be heated up to $kT_2 \approx (3/16)\mu m_{\rm p} v_1^2$, where $\mu \approx 0.6$ is a mean molecular weight and $m_{\rm p}$ is a proton mass, assuming $\gamma = 5/3$.

2.3.3 Characteristic time scale

We summarize here characteristic time-scales that are important indicators to study cluster evolution.

Hubble time: $t_{\rm Hubble}$

The age of the universe is estimated from the Hubble constant:

$$t_{\rm Hubble} \equiv \frac{1}{H_0} = 2.0 \times 10^{10} \text{ yr}, \tag{2.13}$$

where H_0 is taken a value of 50 km s^{-1} Mpc $^{-1}$. This time scale becomes an upper limit on the age of the clusters of galaxies.

Crossing time: $t_{\rm cr}$

According to the hierarchical clustering scenario, we expect the initial state of material in the protocluster to be irregular. Thus the relaxation of the cluster involves the spatial motion of the galaxies, and a lower limit is the crossing time given as

$$t_{\rm cr} \equiv \frac{r}{\sigma_r} \approx 10^9 \left(\frac{\sigma_r}{10^3 \text{km/s}}\right)^{-1} \left(\frac{r}{\text{Mpc}}\right) \text{ yr}, \tag{2.14}$$

where the radial velocity dispersion σ_r has been substituted for the radial velocity v_r.

Two-body relaxation time: $t_{\rm rel}$

Thermodynamic equilibrium could results from elastic collisions between galaxies. The most important elastic two-body collisions in clusters are gravitational. The two-body relaxation time is

$$\begin{aligned} t_{\rm rel} &= \frac{3\sigma_r^3}{4\sqrt{\pi}G^2 m m_{\rm f} n_{\rm f} \ln\Lambda} \\ &\simeq 2\times 10^{10}\text{yr}\left[\left(\frac{\sigma_r}{10^3 \text{km s}^{-1}}\right)^3 \left(\frac{m}{10^{12} M_\odot}\right)^{-1} \left(\frac{m_{\rm f}}{10^{12} M_\odot}\right)^{-1}\right. \\ &\quad \left.\times \left(\frac{n_{\rm f}}{10^3 \text{galaxies Mpc}^{-3}}\right)^{-1} (\ln\Lambda)^{-1}\right] \end{aligned} \tag{2.15}$$

for the relaxation of a galaxy of mass m in a background field of galaxies of mass $m_{\rm f}$ and number density $n_{\rm f}$. Λ is ratio of maximum to minimum impact parameters of collisions which contribute to the relaxation.

Free-Fall time: $t_{\rm ff}$

If we consider heating of the ICM only due to an infall of the primordial gas into the cluster and the compression, the heating time scale is equal to the free-fall time scale given as

$$t_{\rm ff} \approx (G < \rho >)^{-1/2} \approx 3.1 \times 10^9 \left(\frac{D}{1\text{Mpc}}\right)^{3/2} \left(\frac{M_{\rm total}}{10^{14}\ M_\odot}\right)^{-1/2} \text{ yr}, \tag{2.16}$$

where D is the cluster diameter, and $M_{\rm total}$ is the total mass of the cluster.

Cooling time: $t_{\rm cool}$

The primary cooling process for the ICM is radiative cooling due to free-free emission, expressed as

$$t_{\rm cool} = \frac{U}{\epsilon} = \frac{3nkT}{n^2\Lambda(T, Z)} \tag{2.17}$$

where U and ϵ are the internal energy and volume emissivity, and $n = n_{\rm e} = n_{\rm p}$ is the number density in a fully ionized plasma, and Λ is the cooling function. The total emissivity through thermal bremsstrahlung is

$$\epsilon^{ff} \approx 3.0 \times 10^{-27} T^{1/2} n^2 \ {\rm erg\,cm^{-3}s^{-1}}, \tag{2.18}$$

assuming solar abundances and velocity averaged Gount factor of 1.1 in a fully ionized plasma with a temperature of $> 3 \times 10^7$ K (Sarazin 1988, p. 148). Then, the cooling time is calculated as

$$\approx 8.5 \times 10^{10} \left(\frac{n}{10^{-3}\ {\rm cm^{-3}}}\right)^{-1} \left(\frac{T}{10^8\ {\rm K}}\right)^{1/2} \ {\rm yr}. \tag{2.19}$$

Thus the cooling time is longer than the Hubble time for most of the clusters except for the central region.

Ionization equilibration time scale: $t_{\rm ie}$

The time-scale for collisional ionization equilibrium with the plasma calculated by Masai (1994) is

$$t_{\rm ie} \approx 10^{12} n_{\rm e}^{-1} = 3.3 \times 10^8 \left(\frac{n_{\rm e}}{10^{-3}\ {\rm cm^{-3}}}\right)^{-1} \ {\rm yr}. \tag{2.20}$$

Here $n_{\rm e}$ is the electron density.

Equipartition time: $t_{\rm eq}$

If a homogeneous plasma is created in a state in which the particle distribution is non-Maxwellian, elastic collisions will cause it to relax to a Maxwellian distribution on a time scale determined by the mean free paths (Spitzer, 1956, 1978). Electrons will achieve this equipartition on a time scale set roughly by $t_{\rm eq} \equiv \lambda_{\rm e} / < v_{\rm e} >_{rms}$,

$$t_{\rm eq}(e,e) = \frac{3 m_{\rm e}^{1/2} (kT_{\rm e})^{3/2}}{4\pi^{1/2} n_{\rm e} e^4 \ln\Lambda} \approx 3.1 \times 10^5 \left(\frac{\ln\Lambda}{40}\right)^{-1} \left(\frac{T}{10^8 {\rm K}}\right)^{3/2} \left(\frac{n_{\rm e}}{10^{-3}{\rm cm^{-3}}}\right)^{-1} \ {\rm yr}, \tag{2.21}$$

where $T_{\rm e}$ is the electron temperature, and $\ln\Lambda$ is the Coulomb logarithm. The time scale for protons to equilibrate among themselves is $\sqrt{m_{\rm p}/m_{\rm e}} \sim 43$ times longer than the value in $t_{\rm eq}(e,e)$: $t_{\rm eq}(p,p) = 1.3 \times 10^7$ yr. The time scale for the electrons and ions to reach equipartition $T_{\rm e} = T_{\rm i}$ is $t_{\rm eq}(p,e) = (m_{\rm p}/m_{\rm e})t_{\rm eq}(e,e) = 5.7 \times 10^8$ yr.

Conduction time: $t_{\rm cond}$

It is useful to define a conduction time-scale as $t_{\rm cond} \equiv -(d\ln T_{\rm e}/dt)^{-1}$, which can be approximated to $(n_{\rm e}l/\kappa)$ with the scale length of temperature gradient $l \equiv T_{\rm e}/|\nabla T|$ (Sarazin 1988, p.157). Heat conduction is most effective in the cluster core, and $t_{\rm cond}$ increases vary rapidly with radius. The central value to the conduction time scale is given by

$$\begin{aligned} t_{\rm cond}(0) &\approx \frac{n_{\rm e}k}{\kappa}\left(\frac{T}{|\nabla T|}\right)^2 \\ &\approx 5.6 \times 10^7 \left(\frac{n_{\rm p}}{10^{-3}\ {\rm cm}^{-3}}\right)\left(\frac{T_e}{10^8\ {\rm K}}\right)^{-5/2}\left(\frac{\ln\Lambda}{40}\right)\left(\frac{T/|\nabla T|}{0.25\ {\rm Mpc}}\right)^2 {\rm yr}, \quad (2.22) \end{aligned}$$

where $n_{\rm p}$ is the central proton density, and solar abundances have been assumed. Thus heat conduction may be relatively effective in the core of a cluster. At outer radii $r \gtrsim 2r_{\rm c}$, the conduction time is typically 10^2 times longer, and conduction is only marginally effective in the outer parts of the cluster.

Drift time for heavy ions: $t_{\rm drift}$

Heavy ions that diffuse toward the cluster center rapidly reach a drift velocity at which gravitational acceleration is balanced by the slowing effect of collisions (Fabian and Pringle 1977). The time to drift in a small distance dr is then $dt = dr/v_{\rm D}$, where $v_{\rm D}$ is the drift velocity, given as

$$\begin{aligned} t_{\rm drift} \approx\ & 1.2 \times 10^{11}\left(\frac{A}{56}\right)^{-1}\left(\frac{Z}{26}\right)^2\left(\frac{T_{\rm e}}{10^8\ {\rm K}}\right)^{-3/2} \\ & \left(\frac{n_{\rm e}}{10^{-3}\ {\rm cm}^{-3}}\right)\left(\frac{\ln\Lambda}{40}\right)\left(\frac{\sigma_r}{10^3{\rm km/s}}\right)^{-2}\left(\frac{r_{\rm c}}{0.25{\rm Mpc}}\right)^2 {\rm yr}. \quad (2.23) \end{aligned}$$

Rephaeli (1978) has found that the drift time from radii greater than 4 core radii into 1 core radius is always greater than 10^{11} yr the cluster parameters in above equation, with the gas density and temperature evaluated at the cluster center. Since this means the drift time is generally at least ten times the probable age of the cluster, the settling of iron into the cluster center is unlikely to occur. However, this does not mean that the metal abundance is necessarily uniform, because heavy elements might be concentrated in the cluster core if they were injected into the intracluster gas in the center.

Mixing time due to galaxy motions: $t_{\rm mix}$

Motions of galaxies through the intracluster gas may mix the gas. This process has not been treated in any great detail in the literature. It might be reasonable to assume that each galaxy mixes the gas within a wake of radius $R_{\rm W}$. Very roughly, the gas within the whole cluster would be mixed on a time scale given by

$$t_{\rm mix} \approx \frac{1}{n_{\rm gal}\sigma_{\rm r}\pi R_{\rm W}^2}\left(\frac{r_{\rm c}}{R_{\rm W}}\right)^2 , \tag{2.24}$$

where $n_{\rm gal}$ is the number density of galaxies, $\sigma_{\rm r}$ is their velocity dispersion, and $r_{\rm c}$ is the cluster core radius. Depending on whether the galaxy contained any interstellar medium or not, the radius might be ≈ 10 kpc. The resulting value of $t_{\rm mix}$ is generally much longer than $\sim 10^{11}$ yr for any reasonable values of the cluster parameters, and mixing due to galaxy motions is not very important.

Time scale for ram pressure stripping of a galaxy: $t_{\rm rp}$

Sarazin (1979) gave a semi-analytic treatment of the stripping of gas from spherical galaxies, using basically the same formulation as Tarter (1975). Assuming that the ram pressure is much greater than a critical pressure of $2\rho_{\rm ISM}\sigma_*^2$ derived by numerical hydrodynamic simulations, the time-scale for ram pressure stripping of a galaxy, defined as $t_{\rm rp} \equiv (d\ln M_{\rm ISM}/dt)^{-1}$, is

$$\begin{aligned} t_{\rm rp} &\approx \frac{R}{v}\left(\frac{2\rho_{\rm ISM}}{\rho_{\rm g}}\right)^{1/2} \\ &\approx 3\times 10^7 \left(\frac{\rho_{\rm ISM}}{\rho}\right)^{1/2}\left(\frac{v}{10^3{\rm km/s}}\right)^{-1}\left(\frac{R}{20{\rm kpc}}\right) \ {\rm yr}, \end{aligned} \tag{2.25}$$

where $\rho_{\rm ISM}$ is the average interstellar gas density, ρ_g is the intracluster gas density, and v is the galaxy velocity. The calculations of Tarter and Sarazin assume that there are no sources of gas in the galaxy.

2.4 Clusters studied

2.4.1 Selection criteria

For our study, closeness and brightness of a system become much important. Good photon statistics allow ASCA to resolve the temperature structures less than 100 kpc in a whole cluster region. We therefore selected four clusters, Abell 1060 (A1060 for short), AWM 7, the Centaurus cluster and the Perseus cluster with following criteria:

- X-ray bright ($L_{\rm X} \gtrsim 10^{43}$ erg s^{-1})
- nearby ($D_L \lesssim 100$ Mpc which corresponds to $z \lesssim 0.017$)
- rather low temperature ($kT \lesssim 5$ keV)
- axially symmetric X-ray morphology



Figure 2.8: X-ray surface brightness profiles of the 4 clusters from the HRI data (left) and PSPC data (right). The HRI profiles are arbitrarily scaled for comparison between their central features. For the PSPC profiles, the intensity is multiplied by the ratio square of the luminosity distance to the distance of the Perseus cluster to help comparison with the intrinsic luminosity.

General properties and previous X-ray results of the 4 clusters are summarized in Table 2.1. They are all nearby (z =0.0104–0.0183), and X-ray bright clusters with the average ICM temperatures kT =3–6 keV. The clusters, however, are markedly different regarding the strength of the central cool component. Figure 2.8 shows the surface brightness profiles of the 4 clusters from the ROSAT HRI (normalized for comparison) and PSPC data. The Centaurus cluster shows very sharp central peak in the surface brightness profile. The Perseus cluster and AWM7 also show central excess features, but it is not as narrow nor intense as the Centaurus. In contrast, A1060 does not have such a narrow peak, and the size and mass of the cD galaxy is smaller than typical cD galaxies from optical observations. Since the cool component is believed to grow in the course of cluster evolution, the feature of the cool component is a good indicator to examine structure formation of clusters. Their metal distributions also have marked individuality as described in the following subsection.

2.4.2 Abell 1060

A1060 (Hydra I cluster) is the nearest cluster after the Virgo, Fornax, and Centaurus clusters among X-ray luminous ones at a redshift of $z = 0.0114$ (Abell et al. 1989). X-ray emission from this cluster has been observed and studied with *Einstein* (Fitchett and

Table 2.1: General properties for A1060, AWM7, the Centaurus cluster, and the Perseus cluster.

		A1060	AWM7	Centaurus	Perseus
Redshift		0.0114	0.0172	0.0104	0.0183
Luminosity Dist.	Mpc	69	104	63	108
BM/RS[†]		III/C	II?/L?	I-II/cD	II-III/L
Population		103	179	$\gtrsim$ 198	$\gtrsim$ 170
Velocity dispersion	km s^{-1}	676	727	586, 280	$\sim$ 5500
cD galaxy		NGC 3311	NGC 1129	NGC 4696	NGC 1275
R.A.(J2000)[‡]		$10^{h}36^{m}42.6^{s}$	$2^{h}54^{m}27.5^{s}$	$12^{h}48^{m}49.3^{s}$	$3^{h}19^{m}48.2^{s}$
Decl.(J2000)[‡]		$-27°31'42.9''$	$+41°34'42.5''$	$-41°18'40.0''$	$+41°30'42.1''$
l (J2000)	degrees	269.601549	146.336897	302.403583	150.5758384
b (J2000)	degrees	26.486301	−15.635096	21.557991	−13.2611465
$kT_{\rm hot}$ [a]	keV	$3.3^{+0.4}_{-0.3}$	3.6 ± 0.2	3.6 ± 0.4	5.5 ± 0.5
$kT_{\rm hot}$ [b]	keV	3.25 ± 0.05	3.76 ± 0.08	3.75 ± 0.07	6.79 ± 0.11
$L_{\rm X(2-10\ keV)}$ [a]	10^{44} erg s^{-1}	0.25 ± 0.01	1.2 ± 0.07	0.58 ± 0.02	10.0 ± 1.0
$L_{\rm X(0.5-10\ keV)}$ [b]	10^{44} erg s^{-1}	0.41	2.1	0.97	18
Cooling flow rate [c]	$M_\odot$ yr^{-1}	15^{+5}_{-7}	41^{+6}_{-6}	30^{+10}_{-5}	556^{+33}_{-24}
$r_{\rm cool}$ [c]	kpc	79^{+15}_{-26}	103^{+5}_{-8}	81^{+23}_{-18}	185^{+11}_{-11}
$t_{\rm cool}$ [c]	Gyr	$4.7^{+0.4}_{-0.3}$	$1.9^{+0.2}_{-0.2}$	$0.8^{+0.0}_{-0.0}$	$0.9^{+0.0}_{-0.0}$

[†] BM: Bautz-Morgan type, RS: Rood-Sastry type. [‡] Positions of cD galaxy taken from NED. [a] EXOSAT results from Edge and Stewart (1991) [b] ASCA results from Fukazawa et al. (1998) [c] The ROSAT PSPC results from Peres et al. (1998)

Merritt 1988), EXOSAT (Singh et al. 1988), ROSAT, and ASCA (Tamura et al. 1996, 2000). Figure 2.9 shows the PSPC image of the A1060 in the 0.5–2 keV band, the X-ray image is circularly symmetric. The central cool component is very weak or absent in an X-ray spectrum, and cooling flow rate is estimated to be $\dot{M} \lesssim 15 M_\odot$ yr^{-1} (Peres et al. 1998). As shown in figure 2.10, the previous study for the central $r < 20'$ region with ASCA shows that the ICM temperature is quite isothermal at around 3.2 keV and the metal abundance is almost flat at $\sim$ 0.3 solar (Tamura et al. 1996, 2000; Finoguenov et al. 2000).

Richter et al. (1982, 1987) and Hamabe (1993) compiled catalogs of member galaxies of A1060. 103 galaxies have redshifts ranging from 2300 to 5200 km s^{-1}, and can be considered as member galaxies of the cluster. The radial velocity histogram shows that this cluster appears to be isolated in redshift space of 60–70 Mpc. From a Gaussian fit to the radial velocity histogram, the systematic velocity is determined to be 3414 km s^{-1} with an average line-of-sight velocity dispersion of 657 km s^{-1} (Richter et al. 1987). At the center, there are two bright elliptical galaxies NGC 3311($z = 0.0120$) regarded as a cD



Figure 2.9: The PSPC image of the A1060 in the energy range 0.5–2.0 keV.



Figure 2.10: Temperature and abundance distributions of A1060 within a radius of 20′ obtained from ASCA (Tamura et al. 1996).

galaxy, and NGC 3309 ($z = 0.0136$). However, both the size and the mass of NGC 3311 are smaller than the typical cD galaxies. In the southeast near the cluster center, there might be a small foreground group of 3 galaxies from HI observations (McMahon et al. 1992).

2.4.3 AWM7

AWM7 ($z = 0.0172$) is a poor cluster that is not listed in Abell catalog. However, the X-ray luminosity is comparable to other Abell clusters, and it forms part of the Pisces-Perseus supercluster filament at the west of the Perseus cluster. X-ray emission from this cluster has been studied with *Einstein* (Kriss, Cioffi, and Canizares 1983), *Ginga*


G-14). The identification of galaxies in those catalogs is consistent except for one galaxy. NGC 3315 has different positions in RNGC and EUSB. The position of NGC 3315 in RNGC is very close to that of 501 G-47, while 501 G-48 is identified as NGC 3315 in EUSB. However, because the identification of NGC 3315 in EUSB is followed by PGC and RC3, we tentatively adopted the description in EUSB in this study: The parameters for the galaxies 73 and 74 are those for 501 G-47 and 501 G-48, respectively.

Among the program galaxies, 32 are listed in *IRAS* Point Source Catalog (version 2, 1988), and they are indicated by the asterisk (*) following the galaxy name in column (6).

Morphological classifications of the program galaxies given in column (8) were assigned by the examination of the galaxy images on the IIIa-J film copies of the ESO/SERC survey by the author and a student independently with reference to the description given in Sandage (1961). These classifications are in more detail than those in EUSB and bar structure in some galaxies were newly identified.

Membership of the Hydra I cluster given in column (10) is assigned according to the redshifts of the galaxies. For 79 (NGC 3314 was counted twice; see the § 3) among the all program galaxies, the redshifts are available in the literature (Richter et al. 1982; Richter & Huchtmeier 1983, 1987; Palumbo, Tanzella-Nitti, & Vettolani 1983; Fouqué 1987; Richter 1987, 1989; Fairall, Vettolani, & Chincarini 1989; PGC; Huchtmeier & Richter 1989; Fouqué, et al. 1990). The distribution of the redshifts is shown in Figure 1. Among them 74 have redshifts ranging from 2300 to 5190 km s^{-1} and can be considered as members of the cluster (see Richter et al. 1982). They are indicated by "M" in Table 1. Three galaxies (30, 100,

more concentrated to the cluster center than late-type galaxies. The morphology-density relation (Dressler 1980; Postman & Geller 1984; Giovanelli, Haynes, & Chincarini 1986; Binggeli, Tammann, & Sandage 1987) appears to hold for this cluster.

3. MEASUREMENT AND REDUCTION

The galaxy image on the plate was scanned with the PDS 2020GMS microdensitometer at the Kiso Observatory. A square aperture of 17 μm and sampling pitch of 15 μm (1″ on the plate) were adopted for all galaxies. A sufficiently wide area was scanned to determine accurate sky background level. As a result the size of a density image ranges from 517 × 517 pixels

FIG. 1.—Distribution of the redshifts of the program galaxies. The superposed galaxy NGC 3314 was counted twice.

Figure 2.11: *left* – Distribution of the 137 galaxies in A1060 and seven field stars on the UK Schmit plate V3324. Early-type(E-S0/a), late-type(Sa-Irr), and unknown type are denoted by closed circles, open circles, and triangles, respectively. *right* – Distribution of the redshifts of the galaxies (Hamabe 1993).

(Tsuru 1992), ROSAT (Neumann & Böhringer 1995), and ASCA (Xu et al. 1997; Ezawa et al. 1997; Finoguenov & Ponman 1999). As shown in figure 2.12, the X-ray image is elliptically elongated east and west in parallel to the filament structure of Pisces-Perseus supercluster with the ratio to minor to major axis of ~ 0.8. The mapping data of ASCA were previously analyzed by Ezawa et al. (1997). Figure 2.13 shows the radial distribution of temperature, which was found to be fairly flat up to a radius of 40′. They also found a large-scale metallicity gradient. The cool component in the central region is weak and the GIS data can be fit with a single temperature model (Xu et al. 1997).

From optical data, AWM7 was found as a group of galaxies by Albert, White, and Morgan (1977), and was classified as a richness class 0 galaxy cluster (Kriss et al. 1983). For 33 galaxies redshifts are listed in Beers et al. (1984). Koranyi and Geller (2000) have measured 492 redshifts in the direction of the AWM7, and have identified 179 galaxies as member galaxies. Figure 2.14 shows the distribution of the galaxies. The velocity dispersion within 3.6 Mpc is 727 km s^{-1}.

2.4.4 The Centaurus cluster

The Centaurus cluster (Abell 3526, $z = 0.0104$) is the second X-ray bright (next to Perseus cluster) cluster with a dominant cD galaxy, NGC 4696. Figure 2.15 shows the PSPC image of the Centaurus cluster in the 0.5–2 keV band. The Centaurus cluster has been intensively studied in X-rays with *UHURU* (Giacconi et al. 1972), *HEAO 1* (Mitchell and Mushotzky 1980), *Einstein* (Matilsky, Jones, and Forman 1985), EXOSAT



Figure 2.12: The PSPC image of the AWM7 in the energy range 0.5–2.0 keV.



Figure 2.13: Temperature and abundance distributions of AWM7 within a radius of 40′ obtained from ASCA mapping observations (Ezawa et al. 1997).

(Edge and Stewart 1991), *Ginga* (Yamashita 1992), ROSAT (Allen & Fabian 1994), and ASCA (Fukazawa et al. 1994; Ikebe et al. 1999; Churazov et al. 1999). Figure 2.16a shows ASCA results of the emission-measure profiles of the hot and cool component in the central region by Ikebe et al. (1999). The cool component dominates considerably, however, the hot component also exists. The metal abundance profile also shows a central concentration as shown in 2.16b. That significantly implies that the metal distribution accompanies with the cD galaxy.

The presence of the central concentrated cool component suggests that the Centaurus



Figure 2.14: *left* – Position of the sky of all objects with measured redshifts. *right* – Distribution of the redshifts of the galaxies. (Koranyi and Geller 2000)



Figure 2.15: The PSPC image of the Centaurus cluster in the energy range 0.5–2.0 keV.

cluster is a relatively relaxed system. However, some results in both optical and X-rays indicate that there is an evidence of merger. Churazov et al. (1999) shows a two-dimensional map of the gas temperature distribution based on the central 4 pointings with ASCA. Figure 2.17 is the temperature map, and there is a remarkable hot region to south-east. The X-ray emission also shows the excess from the surface brightness model at the region. Furthermore, This hot region corresponds to the position of NGC4709, which is the dominant galaxy of the Cen 45 subcluster group. The member galaxies in



Figure 2.16: (a) Emission-measure profiles of the hot (solid lines) and cool (dashed lines) component, and (b) the iron abundance profile derived from the two-phase model fit. Thick lines show the best-fit models, while thin lines indicate extreme cases at the 90% confidence. (Ikebe et al. 1999).

the Centaurus cluster had been investigated very well by some optical photometric survey. Lucey et al. (1986) discovered a remarkable bimodal velocity distribution in the Centaurus cluster. The two velocity components were found to be roughly centered at about 3000 km s^{-1} and 4500 km s^{-1} and were denoted as Cen30 and cen45, respectively. Figure 2.18 shows the map of member galaxies and the redshift distribution by Stein et al. (1997). The mean velocity and dispersion obtained from the double Gaussian fit are 3397 km s^{-1} and 933 km s^{-1} for Cen30, and 4746 km s^{-1} and 131 km s^{-1} for Cen45. Note that NGC 4696 is at a velocity 2961 $\pm$ 26 km s^{-1}. Thus, these results imply the the smaller Cen 45 group is merging with the dominant Cen 30 group.



Figure 2.17: Color image of the gas temperature distribution derived from the central 4 pointings with ASCA (Churazov et al. 1999).



Figure 2.18: *left* – The sky distribution of the two velocity components of Cen30 and Cen45. *right* – The velocity distributions of the 3° sample with 180 galaxies (Lucey et al. 1986).

2.4.5 The Perseus cluster

The Perseus cluster (Abell 426) is the X-ray brightest cluster of galaxies at a distance of $z = 0.0183$. At the center, the cD galaxy NGC1275 is surrounded by a spectacular low-ionization, emission line nebula (Lynds 1970). The Perseus cluster has been extensively studied with the *UHURU* mission (Gursky et al. 1971; Forman et al. 1972), *Ginga* (Allen et al. 1992), SPARTAN-1 detector (Snyder et al. 1990; Kowalski et al. 1993), ROSAT (Ettori et al. 1998), and ASCA (Fabian et al. 1994; Arnaud et al. 1994; Ezawa et al. 2001). Figure 2.19 shows the PSPC image in the 0.5–2.0 keV band. This cluster forms a part of the Pisces-Perseus supercluster, that includes several other clusters, AWM7, A347, and A262. The elongated direction of the X-ray image is in parallel to the filament structure, in the same way as AWM7. Significant substructure in the PSPC image at east from the cluster center has been discussed as evidence that the cluster is not in a relaxed state and in consistent with a recent merger (Schwarz et al. 1992; Ettori et al. 1998). Ezawa et al. (2001) have examined the temperature and metal abundance distribution based on ASCA mapping observations. Figure 2.4.5 shows the temperature drop in the central region even with the 2-temperature model. The abundance distribution shows not central concentration, but large-scale gentle slope.

Kent and Sargent (1983) identified 167 member galaxies of the Perseus cluster. Figure 2.4.5 shows the distribution of radial velocities fitted with a Gaussian function. The mean velocity is found to be 5470±87 km s−1. It is seen that the velocity dispersion drops from $\sim$ 1300 km s^{-1} at the center to $\sim$ 600 km s^{-1} at the radius of r = 136′. Brunzendorf and Meusinger (1999) also compiled a catalogue of 660 galaxies within a field of about



Figure 2.19: The PSPC image of the Perseus cluster in the energy range 0.5–2.0 keV.



Figure 2.20: Temperature and abundance distributions of Perseus within a radius of 60′ obtained with ASCA (Ezawa et al. 2001). The open and filled circles show the results of the single-temperature and the 2-temperature fit, respectively.

10 square-degrees. In the region of the eastern X-ray excess, they find a significantly higher fraction of spiral and irregular galaxies compared with the morphological mix of the galaxies on the opposite side of the X-ray center.



Figure 2.21: The radial velocity distributions in the Perseus cluster normalized to the local velocity dispersion (Kent and Sargent 1983)

Chapter 3

Instrumentation

This thesis is based on analyzing data that were obtained from observations of X-ray astronomy satellite, ASCA. In this chapter, we briefly summarize the design and performance of each instruments onboard ASCA.

3.1 The ASCA Satellite

ASCA (Advanced Satellite for Cosmology and Astrophysics) is the forth Japanese X-ray astronomy satellite (Tanaka et al. 1994) following Hakucho (1979), Tenma (1983) and Ginga (1987). ASCA was launched on February 20, 1993 using the three-stage solid-propellant M-3SII-7 rocket from Kagoshima Space Center of ISAS, and achieved a near-circular orbit with a perigee of 520 km, an apogee of 625 km, and an inclination of 31°. Figure 3.1 shows the in-orbit configuration of ASCA. It weighs 420 kg, and has an octagonal-shaped body of a diameter 1.2 m where six solar panels are attached. ASCA re-entered the atmosphere on March 2, 2001, the day after Prof. Minoru Oda passed away.

ASCA is equipped with four X-ray telescopes (XRTs) at the top of an extendible optical bench. X-rays that the XRTs correct with the large effective areas are focused on four position-sensitive X-ray detectors; two SIS (Solid-state Imaging Spectrometer) detectors called SIS0 and SIS1, and two GIS (Gas Imaging Spectrometer) detectors called GIS2 and GIS3.

3.2 X-ray Telescope (XRT)

The XRT onboard ASCA has enabled cosmic X-ray imaging studies above energies of a few keV up to 10 keV for the first time (Serlemitsos et al. 1995). Soft X-rays are totally reflected off a smooth surface, when their incident angle measured from the surface is shallower than a certain critical value. This phenomenon is known as grazing-incidence reflection. The critical angle, typically of order 1 degree or so, is inversely proportional



Figure 3.1: In orbit configuration of the ASCA satellite

Table 3.1: Design parameters and performance of the ASCA XRT.

Focal length	3500 mm
Incident angle	$0.24^\circ - 0.7^\circ$
Geometrical area	558 cm^2 / telescope
Field of view	$\sim 24'$ (FWHM at 1 keV) / $\sim 16'$ (FWHM at 7 keV)
Energy range	$\lesssim$ 10 keV
Effective area of four XRTs	$\sim$ 1300 cm^2 (1 keV) / $\sim$ 600 cm^2 (7 keV)
Half power diameter	$\sim$ 3 arcmin

to the X-ray energy and increases with increasing free electron density of the reflecting material. X-ray telescope mirrors use this mechanism usually in so-called Wolter type I configuration, which employs paraboloid and hyperboloid surfaces as the primary and the secondary mirrors so as to remove the first-order aberration. The two mirrors have a common focus and reflect X-rays in series (double reflection).

The reflector shells of the ASCA XRT are all made of thin (127 μm) aluminum foils, bent in a conical form to approximate paraboloid or hyperboloid. The bent foils are $\sim$ 10 μm lacquer-coated to improve the surface smoothness, and then $\sim$ 50 nm gold-evaporated to increase the reflectivity. The 120 of these foils are closely packed together in an onion-ring configuration with a typical spacing of 1 mm, to cover a large fraction of the mirror aperture. The innermost and the outermost shells have diameters of 120 mm and 345 mm, respectively. In practice, the foils are produced and packed in four quadrants, and aligned by 13 alignment bars into 14 sectors. The first and the last sector is masked because shape error is severe. Four quadrants make up the "paraboloid" section of 100 mm long, and another set of four quadrants make up the "hyperboloid" section. One XRT unit thus fabricated weighs about 9.8 kg. Design parameters and performance of the XRT are shown in table 3.1.



Figure 3.2: Effective area of the ASCA XRT (4 XRTs summed up, on-axis) compared with that of polished mirrors used in previous missions.

The XRT has successfully achieved a very large effective area and a wide energy range, as shown in figure 3.2. When the four units are summed up, the effective area reaches $\sim$ 1100 cm^2 below $\sim$ 2 keV, almost by a factor of 6 and 2 larger than those of Einstein and ROSAT respectively. A sudden drop in the reflectivity at 2.2 keV is caused by the M-edge of the gold used as reflective coating. The XRT maintains a very large reflecting area towards higher energies, e.g. $\sim$ 600 cm^2 at 7 keV, with the upper energy bound as high as $\sim$ 10 keV. X-ray imagery above 3–4 keV has never been achieved by polished mirrors used in previous missions.

Although the XRT has a large effective area even at $\sim$ 10 keV, it sacrifices the imaging quality: large-extended point spread function, and stray light which comes from outside of the field of view caused by abnormal reflection. These effects on our analysis are described in section 4.1.

3.3 Solid-state Imaging Spectrometer (SIS)

The SIS experiment is the first X-ray sensitive charge coupled devices (CCDs) detector system in orbit. Figure 3.3 shows a cross section view of the SIS camera. The SIS system consists of two detectors, SIS0 and SIS1, and each SIS detector is made up of four CCD chips of 11 mm square. The 4 CCD chips are aligned in square with narrow gaps to cover about $20' \times 20'$ square region on the sky. Each chip has 4096 by 4096 pixels of 27 μm square each, and a depletion layer of about 40 μm thick which ensures an improved efficiency for harder X-rays than conventional CCDs. Design parameters and performance of the SIS in brief are summarized in table 3.2.

Figure 3.4a illustrates the quantum efficiency of the SIS as a function of incoming X-ray energy. Thus the SIS sensitivity covers approximately 0.4–10 keV. The CCD chips and preamplifiers are cooled down to -60 °C with a thermo-electric cooler (TEC) from the backside of the chips in order to reduce thermal noise down to $N \sim 5$ electrons level.



Figure 3.3: Cross section view of the SIS camera

Table 3.2: Design parameters and performance of the SIS

Energy Band	0.4–12 keV
Energy Resolution	2% at 5.9 keV (FWHM)
Effective Area	11×11 mm square per chip
Optical Blocking Filter	100 nm Lexan film coated with 40 nm aluminum
Number of pixels in Image Region	420 pixels × 422 lines per chip
Pixel Size	27 μm
Thickness of Depletion Layer	$\sim$ 40 μm

Thus the SIS achieves an energy resolution of about 150 eV FWHM over the whole energy range (figure 3.4b); this is the best energy resolution ever achieved by non-dispersive X-ray spectrometers so far put into orbit.

3.4 Gas Imaging Spectrometer (GIS)

The GIS is the imaging gas scintillation proportional counter (Makishima et al 1996; Ohashi et al. 1996). The application of the gas scintillator technique to the Japanese X-ray satellite is the second attempt after the successful operation of the SPC experiment on board Tenma (Tanaka et al. 1984; Koyama et al. 1984). The GIS is a general-purpose X-ray imaging spectroscopy system. It consists of the two detector assemblies, namely



Figure 3.4: (a) Detection efficiency of the SIS as a function of incoming X-ray energy. K-edges of O (0.53keV), Al (1.56keV), and Si (1.84keV) are clearly seen in the figure. This efficiency does not includes optical blocking filter. (b) Energy resolutions of the SIS as a function of incoming X-ray energy for the single event. Energy resolutions with different read-out noise N are plotted ($N \sim 5$ for the SIS). The read-out noise levels are given as the equivalent number of electrons.

GIS2 and GIS3 serving as X-ray detectors. The GIS2 and GIS3 are coupled to two of the four XRTs, and measure pulse-heights and positions of X-rays reflected by the XRTs, photon-by-photon basis.

Figure 3.5 shows a schematic view of the GIS sensor system. X-rays reflected by the XRT enter through the window, whose electric potential is held at -6000 V, and are absorbed in the drift region. Through photo-ionization, primary electrons are generated on average at the rate of one electron per 21.5 eV. The electron cloud thus created slowly drifts to the intermediate mesh (-5300 keV), and then the cloud is accelerated due to the strong field toward the ground mesh which is placed in front of the quartz window. In this process, the electrons excite Xe and produce a large number of UV photons of $\sim$ 170 nm wavelength. The excitation energy for one UV photon is $\sim$ 10 eV. Through the quartz window, these UV photons are collected by the imaging photo-multiplier tube (IPMT) which measures light distribution and the overall intensity of UV flux, which is proportional to the X-ray energy with accuracy of several %. We can obtain the position information from the anode signals, and derive the pulse-hight (PH) and the rise-time (RT) information from the last dynode. Design parameters and performance of the GIS are summarized in table 3.3, and detection efficiency and energy resolution are shown in figure 3.6.

There are a number of monitor data in the GIS system to support monitoring its proper function and to help estimation such as background and dead time (Ohashi et al. 1996). In the PH vs. RT plane, there are 6 scalers which measure the monitor counts (see figure 3.7). The upper PH discriminator sets one threshold line, and lower and upper